

DIATOM ASSEMBLAGES FROM GOSHIKI-NUMA

Richard W. Jordan & Hayato Abe

Department of Earth & Environmental Sciences

As with the larger lakes (Lakes Hibara, Onogawa and Akimoto) on the Urabandai Plateau, Goshiki-numa's ponds and lakes were formed in 1888 following the last eruption of Mt. Bandai. However, unlike these larger circum-neutral lakes, many of the ponds within the Goshiki-numa system are weakly to strongly acidic (pH 3-6), although some are >pH7 (e.g. Lake Yanagi-numa). It is widely accepted that these acidic ponds receive their water supply from underneath Mt. Bandai itself, whereas the larger lakes are fed by the mountains to the north. Historical studies of the water characteristics of Goshiki-numa's ponds and lakes are relatively few, but enough information exists to be able to reconstruct the pH changes over the last 70 years (Table I).

Location	Sampling date												
	① July 1931	② 16-20 Aug. 1935	③ 8 Sept. 1937	④ Aug. 1940	⑤ 13-16 Oct. 1952	⑥ 24 July 1964	⑦ 6-8 Aug. 1968	⑧ Aug. 1969	⑨ June 1980	⑩ Oct. 1986	⑪ 1988?	⑫ 1996?	⑬ June-Oct. 1996
Lake Aka-numa	4.1	3.3	3.0	3.5	4.0	3.9	3.9					3.4	
Lake Benten-numa	4.9	4.5	4.7	4.5	4.8	4.4	4.7					4.0-4.5	
Lake Ruri-numa	4.5	4.3	4.2	4.7	4.8	4.2	4.3				4.44	4.0	
Lake Midoro-numa		5.8	6.2	6.3	6.0	4.6	6.9					5.7	
Lake Ao-numa		4.4		4.7	4.8	4.1	4.2					6.5	
Lake Bishamon-numa	5.0	4.5-4.7	6.2	5.0	6.5	4.8	6.2		5.8-6.4	6.5	6.44		5.6-6.3
Lake Tatsu-numa		6.2	6.4		5.0	5.0-5.4		6.7				6.0	
Lake Yaroku-numa		6.5	6.6*	6.7	6.8	6.6	6.9						
Lake Yanagi-numa		7.2	7.1	7.3	6.9	5.3		6.7					

*6.3 in Yoshimura *et al.* (1936a)

Table I. Historical pH data from the Japanese literature on some of the Goshiki-numa lakes. References 1-3 were taken from Yoshimura *et al.* (1937), although Ref.1 (Yoshimura, 1932) and Ref.2 (Yoshimura *et al.*, 1936a) were the original papers. Other references include Negoro (1944; Ref.4), Fukushima (1964; Ref.5), Tanabe *et al.* (1964; Ref.6), Kato & Aita (1970; Refs 7 and 8), Tanaka (1992; Refs 9 and 10), Chiba (1988; Ref.11), Yoshitake & Fukushima (1996; Ref.12), and Sugawara *et al.* (1996; Ref.13).

Some of the earliest studies were conducted by Yoshimura (1932, 1935) and Yoshimura *et al.* (1936a, 1936b, 1937) who made limnological surveys of the lakes and ponds between 1931-1937. Later investigations were made by Negoro (1942, 1944, 1968, 1970), Fukushima (1964), Tanabe *et al.* (1964), Adachi (1969, 1970), Kato & Aita (1970), the JIBP-PF Research Group of acid lakes in Urabandai (1975), and Yoshitake & Fukushima (1996). More recently, detailed information on Lake Bishamon-numa has been reported by Sugawara *et al.* (1999). Many of these studies reported on the diatom species found in each lake. Here, a summary of the important species is presented in Table II.

Over the last 3 years a number of water and surface sediment samples have been collected from the various lakes within the Goshiki-numa National Park (Fig.1; Table III). The diatom assemblages within these samples have now been identified, photographed, counted, and compared with the physico-chemical data set (temperature, salinity, conductivity, pH, Fe^{2+} , Fe^{3+}) of each waterbody. The sample preparation methods have been discussed previously. Most of our physico-chemical measurements fall within the ranges of those taken by previous researchers. The dominant species found in each lake are given in Table III. In addition, two short gravity cores were taken in Lakes Bishamon-numa (5-7-1996; Fig.2) and Benten-numa (29-9-1997; Fig.3). The diatom assemblages within the Bishamon-numa core were previously reported by Shiono & Jordan (1997), but the data has been replotted here in order to compare the assemblages with those of the Lake Benten-numa core. The short cores have not been dated yet, but are presumed to cover most of the post-eruption period (1888-1996/7).

The historical pH data set for Lake Bishamon-numa (Table I) together with our pH values (Table III) show that the lake pH has moderately fluctuated over the last 70 years (pH 4.5-6.8), but appears to have been more stable (pH 5.6-6.8) since the late 1960's. The diatom assemblages (Fig.2) do not show any major fluctuations throughout most of the core, however, in the top part of the core (samples 14-16) a change occurred, characterized by a sharp increase in the percentage of *Aulacoseira ambigua* (called *Aulacoseira* sp. in Shiono & Jordan, 1995; 1997). At present we are not sure why this change occurred, but it may be related to the shift in pH towards more circum-neutral

Location	Important diatom species			
	16~20 August 1935 Yoshimura <i>et al.</i> (1936)	August 1940 Negoro (1942)	August 1940 Negoro (1944)	13~16 October 1952 Fukushima (1964)
Lake Aka-numa		<i>Eunotia</i> sp. (= <i>E. levistriata</i> ?)	<i>Eunotia septentrionalis</i> var. <i>intermedia</i> , <i>E. s.</i> var. <i>levistriata</i> , <i>Pinnularia</i> <i>braunii</i> var. <i>amphicephala</i> *	
Lake Benten-numa	<i>Asterionella</i> , <i>Melosira</i> ‡, <i>Navicula</i>		<i>Asterionella formosa</i>	<i>Rhopalodia gibberula</i>
Lake Ruri-numa	<i>Asterionella</i> , <i>Epithemia</i> , <i>Fragilaria</i>		<i>Asterionella formosa</i>	
Lake Midoro-numa	<i>Asterionella</i> , <i>Melosira</i> ‡, <i>Diatoma</i>			<i>Pinnularia subcapitata</i>
Lake Ao-numa	<i>Asterionella</i> , <i>Diatoma</i> , <i>Surirella</i>		<i>Asterionella formosa</i>	
Lake Bishamon-numa	<i>Asterionella</i>		<i>Pinnularia braunii</i> var. <i>amphicephala</i> *, <i>Asterionella</i> <i>formosa</i>	
Lake Tatsu-numa	<i>Asterionella</i> , <i>Melosira</i> ‡, <i>Surirella</i>			<i>Rhopalodia gibba</i>
Lake Yaroku-numa	<i>Asterionella</i> , <i>Tabellaria</i> , <i>Fragilaria</i> , <i>Diatoma</i>			<i>Fragilaria</i> sp., <i>Rhopalodia gibba</i>
Lake Yanagi-numa	<i>Asterionella</i> , <i>Melosira</i> ‡, <i>Navicula</i> , <i>Fragilaria</i>			<i>Fragilaria</i> sp., <i>Rhopalodia</i> <i>gibba</i> , <i>Synedra ulna</i> var. <i>biceps</i> , <i>Mastogloia smithii</i> , <i>Epithemia turgida</i>

‡ These *Melosira* spp. are now probably *Aulacoseira* spp.

* This is probably a misidentification, and may be what is now called *P. acidojaponica* (Idei & Mayama 2001)

Location	Important diatom species		
	Negoro (1970)	JIBP-PF (1975)	Yoshitake & Fukushima (1996)
Lake Aka-numa			<i>Nitzschia capitellata</i>
Lake Benten-numa			<i>Eunotia exigua</i> , <i>Surirella</i> sp., <i>Nitzschia capitellata</i>
Lake Ruri-numa			<i>Eunotia exigua</i>
Lake Midoro-numa			<i>Aulacoseira</i> sp., <i>Melosira</i> <i>varians</i> , <i>Eunotia exigua</i>
Lake Ao-numa			<i>Fragilaria vaucheriae</i> var. <i>capitellata</i> , <i>Anomoeoneis</i> <i>neoxilis</i> , <i>Nitzschia obtusa</i> var. <i>scalpelliformis</i>
Lake Bishamon-numa			
Lake Tatsu-numa	<i>Melosira italica</i> ‡, <i>Surirella</i> <i>tenera</i> , <i>Nitzschia obtusa</i> , <i>Fragilaria construens</i> var. <i>venter</i> , <i>Gomphonema gracile</i>	<i>Melosira italica</i> ‡, <i>Melosira</i> <i>granulata</i> ‡, <i>Nitzschia obtusa</i> var. <i>scalpelliformis</i> , <i>Nitzschia</i> <i>obtusa</i> var. <i>lapidula</i>	<i>Fragilaria</i> sp.
Lake Yaroku-numa			
Lake Yanagi-numa			

‡ These *Melosira* spp. are now *Aulacoseira* spp.

Table II. Historical diatom data from the Japanese literature on some of the Goshiki-numa lakes.




Figure 1. Map showing the water and core sampling point locations in the Goshiki-numa area.

Location	Sampling date	Temperature (°C)	pH	Dominant diatom species
Lake Aka-numa	1-10-1998	6.5-7.8	3.9	<i>Eunotia exigua</i> and <i>Pinnularia abaujensis</i>
	21-9-2000	14.2-14.8	3.8-4.1	
Lake Benten-numa	1-9-1997	15.2	4.8	<i>Surirella linearis</i> var. <i>linearis</i> and <i>Surirella</i> sp.1
	21-9-2000	22.3-23.1	4.1-4.5	
Lake Ruri-numa	1-9-1997	15.2	4.6	<i>Fragilaria</i> spp., <i>Brachysira brebissonii</i> and <i>Eunotia arcus</i>
	21-9-2000	21.3-22.1	4.5	
Lake Midoro-numa	1-9-1997	11.6	5.3	<i>Eunotia arcus</i>
	1-10-1998	11.3-12.0	5.4	
	21-9-2000	17.3-18.1	5.4	
Lake Ao-numa	1-10-1998	8.8-9.3	5.7	<i>Gomphonema</i> sp.
	21-9-2000	20.8-21.6	6.6	
Lake Bishamon-numa	1-9-1997	15.6	6.8	<i>Fragilaria</i> spp. and <i>Aulacoseira italica</i>
	1-10-1998	9.6-9.7	6.8	
	21-9-2000	21.6-22.7	5.9-6.2	
Lake Tatsu-numa	1-9-1997	13.4	7.2	<i>Fragilaria</i> spp. and <i>Aulacoseira italica</i>
	1-10-1998	10.3-11.2	7.0	
	21-9-2000	17.6-18.2	6.4-6.9	
Lake Yaroku-numa	21-9-2000	20.8-21.7	6.8-7.2	<i>Fragilaria</i> spp., <i>Cocconeis placentula</i> and <i>Gomphonema</i> spp.
Lake Yanagi-numa	1-9-1997	14.5	7.3	<i>Fragilaria</i> spp., <i>Rhoicosphenia abbreviata</i> and <i>Epithemia adnata</i>
	1-10-1998	9.2-10.0	7.2	
	21-9-2000	19.3-19.9	6.6-6.9	

Table III. Physico-chemical and diatom data of some of the Goshiki-numa lakes collected during the present study.

conditions. Increases in *A. ambigua* percentages were also seen in Lake Hibara and Lake Onogawa core tops (Shiono & Jordan, 1997).

The historical pH data set for Lake Benten-numa (Table I) together with our pH values (Table III) show that the lake pH has hardly changed over the last 70 years. The diatom assemblages within our core (Fig.3) also reflect this general lack of pH change, however, at the bottom of the core (samples 01-03) a different flora existed. One can only speculate that these samples correspond to a deposition time before the historical data set began (i.e. prior to 1931). This early diatom assemblage is dominated by attached and benthic diatoms, whilst later samples contain high percentages of *Aulacoseira ambigua* and *A. italica*, both planktonic forms. This suggests that Lake Benten-numa was much shallower in the earlier years (i.e. shallower than the 6.7m reported by Yoshimura *et al.*, 1936a). The planktonic diatom assemblage of Lake Benten-numa in the summer of 1935 (Yoshimura *et al.*, 1936b) was dominated by *Asterionella*, whilst *Aulacoseira* (cited as *Melosira*) and *Navicula* (both important genera in our core) were rare. However, this could be interpreted as unrepresentative of the spring diatom bloom, which by August would have sunk out of the water column and would have been lying on the sediment surface.

In the future it is hoped that further limnological information can be gathered from Goshiki-numa and that more short cores can be taken and possibly dated. In this way it may be possible to design a local proxy-pH meter using the pH tolerances of living diatom assemblages, and to reconstruct past pH changes using the change in diatom assemblages downcore. At present the pH range of Goshiki-numa lakes lies between pH 3.8-7.3, however, the recent volcanic tremors on Mount Bandai (article in Japan Times, 1st January 2001) may indicate that the volcano is close to a small-scale eruption. If that happens, it may cause a decrease in the pH value of the source waters, and thus eventually of Goshiki-numa's lakes. In order to develop the proxy-pH meter our present range needs to be extended by incorporating Lakes Akadoro-numa, Midori-numa and Mosen-numa (pH3.2-3.9) and the fumaroles near Nakanoyu Spa (JIBP-PF Research Group of acid lakes in Urabandai, 1975). A survey of acidobiontic diatom communities has already been carried out elsewhere (Lake Okama, Zao Spa, and Lake Katanuma) by our group, where it was shown that the diatom assemblages living in waters of pH1-3

Figure 2. The downcore changes in diatom assemblage in a Bishamon-numa core. Sample 16 represents the top of the core.





Figure 3. The downcore changes in diatom assemblage in a Benten-numa core. Sample 23 represents the top of the core.



are more specialized and less diverse (basically dominated by *Pinnularia* spp.) than those found in waters >pH3 (Jordan, 2001). In addition, the area around Banso Spa (pH 8.3; Chiba 1988), close to Lake Hibara, may provide us with slightly alkaline samples.

References

- Adachi, R. (1969). Phytoplankton of Lake Tatsu-numa. In: *Report of JIBP-PF in 1968 – Biological productivity of disharmonic lakes*. pp.53-56. [in Japanese]
- Adachi, R. (1970). The study of phytoplankton in Lake Tatsu-numa. The important species and its monthly change. In: *Report of JIBP-PF in 1969 – Biological productivity of disharmonic lakes*. pp.20-25. [in Japanese]
- Chiba, S. (1988). Water quality of Lake Inawashiro and lakes in Ura-bandai district. In: *Bandai-san – centenary of the 1888 eruption of Bandai-san. Journal of Geography* 97: 134-139.
- Fukushima, H. (1964). Diatoms from Bandai Goshikinuma lake group, Central Japan. *Bulletin of Yokohama Municipal University, Natural Science* 16: 55-75.
- Idei, M. & Mayama, S. (2001). *Pinnularia acidojaponica* M. Idei et H. Kobayasi sp. nov. and *Pinnularia valdetolerans* Mayama et H. Kobayasi sp. nov. – new diatoms from Japanese extreme environments. In: *Lange-Bertalot-Festschrift* (Jahn, R., Kociolek, J.P., Witkowski, A. & Compère, P., eds), Gantner, Ruggell. (in press).
- JIBP-PF Research Group of acid lakes in Urabandai (1975). Productivity of the community of acidic lakes in Urabandai, with special reference to Lake Tatsu-numa, one of the disharmonic lakes. In: *Productivity of communities in Japanese inland waters* (Mori, S. & Yamamoto, G., eds), pp.150-197. University of Tokyo Press.
- Jordan, R.W. (2001). Taxonomy, morphology and distribution of two *Pinnularia* species from acidic lakes and rivers in Yamagata and Miyagi Prefectures, Northeast Japan. In: *Lange-Bertalot-Festschrift* (Jahn, R., Kociolek, J.P., Witkowski, A. & Compère, P., eds), Gantner, Ruggell. (in press).
- Jordan, R.W. & Abe, H. (1997). A brief review of the diatom flora of Goshiki-numa and details of a new sampling programme. *The Report of the Urabandai Limnological Station, Faculty of Science, Yamagata University* No.5: 3-6.
- Kashimura, T. (1988). Vegetation in and around Lakes Goshikinuma. In: *Bandai-san – centenary of the 1888 eruption of Bandai-san. Journal of Geography* 97: 134-139.
- Kato, T. & Aita, N. (1970). Geochemical investigations of the lakes in Ura-bandai district. In: *Report of JIBP-PF in 1969 – Biological productivity of disharmonic lakes*. pp.53-59. [in Japanese]
- Negoro, K. (1942). Studies on the vegetation of the mineralogenous acidotrophic waters of Japan in 1940. *Jour. Plantecol.*, 2: 25-34.
- Negoro, K. (1944). Untersuchungen über die Vegetation der mineralogen-azidotrophen Gewässer Japans. *Sci. Rep. Tokyo Bunrika Daigaku*, Sec. B, 6 (101): 231-374.
- Negoro, K. (1968). The algae of Bandai volcano's Lake Akanuma, and the plankton of Goshikinuma's Lake Yanaginuma. In: *Report of JIBP-PF in 1967 – Biological productivity of disharmonic lakes*. pp.59-61. [in Japanese]
- Negoro, K. (1970). Littoral sessile algae of Lake Tatsu-numa. In: *Report of JIBP-PF in 1969 – Biological productivity of disharmonic lakes*. pp.18-19. [in Japanese]
- Shiono, M. & Jordan, R.W. (1995). Recent diatoms of Lake Hibara, Fukushima Prefecture. *Diatom* 11: 31-63.
- Shiono, M. & Jordan, R.W. (1997). Recent diatom history of four lakes formed by the eruption of Mount

- Bandai. *The Report of the Urabandai Limnological Station, Faculty of Science, Yamagata University* No.5: 27-34.
- Sugawara, K., Hino, S., Nakazato, R., Ochiai, M. & Satoh, Y. (1999). Physicochemical and biological characteristics of Lake Bishamon-numa in Urabandai, Fukushima Prefecture. *The Japanese Journal of Limnology* 60: 367-377.
- Tanabe, K., Kanazawa, T. and six students (1964). pH values of water in the lakes of Goshiki-numa Group, Fukushima Prefecture. *Tohoku-Chiri* 16: 205.
- Tanaka, M. (1992). The lakes in Japan. Nagoya University Press. [in Japanese]
- Yoshimura, S. (1932). Preliminary study on the local limnology of volcanic lakes around Volcano Bandai. *Chirigaku-Hyoron* 8: 46-66. [in Japanese]
- Yoshimura, S. (1935). Lake Akanuma a siderotrophic lake, at the foot of volcano Bandai, Fukushima Prefecture, Japan. *Proc. Imp. Acad. Tokyo* 11, 10: 426-428.
- Yoshimura, S., Negoro, K. & Yamamoto, S. (1936a). Limnological reconnaissance of the Goshikinuma Lake Group of Volcano Bandai, Fukushima Prefecture (I). *Chirigaku-Hyoron* 12: 1-17. [in Japanese]
- Yoshimura, S., Negoro, K. & Yamamoto, S. (1936b). Limnological reconnaissance of the Goshikinuma Lake Group of Volcano Bandai, Fukushima Prefecture (II). *Chirigaku-Hyoron* 12: 42-69. [in Japanese]
- Yoshimura, S., Negoro, K. & Yamamoto, S. (1937). Limnological reconnaissance of the Goshikinuma Lake Group of Volcano Bandai, Fukushima Prefecture (III). *Chirigaku-Hyoron* 13: 1147-1152. [in Japanese]
- Yoshitake, S. & Fukushima, H. (1996). Distribution of living diatom communities in inorganic acid waters in Japan. *Abstracts of the 14th International Diatom Symposium, September 2-8, 1996, Tokyo*. p.133.

3. 研究実績

この章に収録した論文は、いずれ学術雑誌に原著等として発表される予定です。
特に引用を希望される方は、引用の可否について下記へお問い合わせ下さい。

問い合わせ先

名前：原 慶明

住所：990-8560 山形市小白川町1-4-12 山形大学理学部生物学科

電話：023-628-4610

Fax：023-628-4625

e-mail:hara@sci.kj.yamagata-u.ac.jp